	Riverhe	ad 2020 - Tick Infection Ra	ites	
Collection site is located within the 11949 zip code				
Infection Rate (%)	Pathogen	Disease	Stage	Tick Species
38.0	B. burgdorferi	Lyme Disease	Adult	Blacklegged Tick
30.0	B. microti	Babesiosis	Nymph	Blacklegged Tick
22.0	B. burgdorferi	Lyme Disease	Nymph	Blacklegged Tick
14.0	A. phagocytophilum	Anaplasmosis	Nymph	Blacklegged Tick
10.0	B. microti	Babesiosis	Adult	Blacklegged Tick
4.0	A. phagocytophilum	Anaplasmosis	Adult	Blacklegged Tick
4.0	B. miyamotoi	Relapsing Fever	Adult	Blacklegged Tick
4.0	E. ewingii	Ehrlichiosis	Adult	Lone Star Tick
2.0	E. chaffeensis	Ehrlichiosis	Adult	Lone Star Tick
2.0	E. ewingii	Ehrlichiosis	Nymph	Lone Star Tick
2.0	B. miyamotoi	Relapsing Fever	Nymph	Blacklegged Tick
2.0	Powassan virus	Powassan Virus Disease	Adult	Blacklegged Tick
0.0	E. chaffeensis	Ehrlichiosis	Nymph	Lone Star Tick
0.0	Powassan virus	Powassan Virus Disease	Nymph	Blacklegged Tick

⁻While nymphal ticks often have lower infection rates than adult ticks, they are typcially responsible for causing more disease in humans, as their smaller size allows them to avoid notice